



Clark County School District
**Early Childhood Special Education (ECSE)/
Pre-Kindergarten (Pre-K) Immunization Requirements
2025–2026 School Year**

Rev. 06/18/2025

Below are the immunizations, per Nevada Revised Statutes (NRS) 392.435, which are required before entering an ECSE/Pre-K Program. Once enrolled, a child who turns 4 years of age has 30 days to complete the required immunizations pursuant to NRS 392.435 and 394.192.

***See the Clark County School District Health Services Department K–12 Quick Immunization Reference Guide 2025–2026 School Year for interval dosing and additional guidelines.

<u>AGE 3</u>	<u>AGE 4</u>
<u>4 DTP or DTaP (lab titer NOT acceptable)</u> (minimum age: 6 weeks)	<u>5 DTP or DTaP (lab titer NOT acceptable)</u> <ul style="list-style-type: none">• 5th dose not needed if 4th dose given on or after the 4th birthday and at least 6 months after the previous dose.
<u>2 HEP A (lab titer acceptable)</u> (minimum age: 12 months)	<u>2 HEP A (lab titer acceptable)</u> <ul style="list-style-type: none">• 2nd dose must be given at least 6 months after the 1st dose.
<u>3 HEP B (lab titer acceptable)</u> (minimum age: Birth)	<u>3 HEP B (lab titer acceptable)</u> <ul style="list-style-type: none">• 1st and 2nd dose must be separated by 4 weeks.• 2nd and 3rd dose must be separated by at least 8 weeks.• 3rd dose at least 16 weeks after the 1st dose. (Minimum age for final dose [3rd or 4th dose] is 24 weeks of age.)
<u>1 MMR (lab titer acceptable)</u> (minimum age: 12 months)	<u>2 MMR (lab titers acceptable)</u> <ul style="list-style-type: none">• Compliant if 2 doses at least 4 weeks apart.
<u>3 Polio (lab titer NOT acceptable)</u> (minimum age: 6 weeks)	<u>3-4 Polio (lab titer NOT acceptable)</u> <ul style="list-style-type: none">• 1st and 2nd dose, and 3rd dose must be at least 4 weeks apart.• Final dose (3rd or 4th dose) at least 6 months after previous dose and on or after 4th birthday.• 4th dose not needed if 3rd dose is at least 6 months from the previous dose and on or after 4th birthday.
<u>1 Varicella (lab titer acceptable)</u> (minimum age: 12 months)	<u>2 Varicella</u> <ul style="list-style-type: none">• 1st and 2nd dose should be separated by 3 months. Compliant if at least 4 weeks apart.